

AIRX[®] 78 PLUS

Disinfectant Cleaner
& Odor Counteractant

Efficacy
Data

EPA REG. # 10324-155-44089

Efficacy

Hospital Disinfection (at 1 ounce per gallon)

AIRX78 PLUS is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum and 400 ppm hard water at 1 ounce of this product per gallon of water (660 ppm active). Treated surfaces must remain wet for 10 minutes. (Testing is performed per the AOAC UDT/GST method (DIS/TSS-1). Sixty carriers are required on 3 separate lots, one of which must be > 60 days old against *Pseudomonas aeruginosa*, *Salmonella enterica* and *Staphylococcus aureus*. Killing of 59 out of 60 carriers is required (total carriers = 540).)

| ORGANISM | CARRIER POPULATION | SAMPLE | # CARRIERS | # POSITIVE |
|---|------------------------------------|---------------|------------|------------|
| <i>Pseudomonas aeruginosa</i> ATCC #15442 | 3.9 X 10 ⁴ CFU/Carrier | A | 60 | 0/60 |
| | | (60 Days Old) | | |
| | | B | 60 | 0/60 |
| <i>Salmonella enterica</i> ATCC #10708 | 1.03 X 10 ⁶ CFU/Carrier | C | 60 | 1/60 |
| | | (60 Days Old) | | |
| | | A | 60 | 1/60 |
| <i>Staphylococcus aureus</i> ATCC #6538 | 7.0 X 10 ⁴ CFU/Carrier | B | 60 | 1/60 |
| | | (60 Days Old) | | |
| | | C | 60 | 0/60 |
| | | A | 60 | 0/60 |
| | | (60 Days Old) | | |
| | | B | 60 | 0/60 |
| | | C | 60 | 0/60 |
| | | (60 Days Old) | | |
| | | A | 60 | 0/60 |
| | | B | 60 | 0/60 |
| | | (60 Days Old) | | |
| | | C | 60 | 0/60 |

Supplemental Organisms

(Testing is performed per the AOAC UDT/GST method. Ten carriers are required on 2 separate lots against each supplemental organism. Killing of 10 out of 10 carriers is required (total carriers = 20).)

| ORGANISM | CARRIER POPULATION | SAMPLE | # CARRIERS | # POSITIVE |
|--|------------------------------------|--------|------------|------------|
| <i>Acinetobacter baumannii</i> ATCC 19003 | 5.1 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Acinetobacter Iwoffii</i> ATCC 15309 | 5.7 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Bordetella bronchiseptica</i> ATCC 10580 | 9.4 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Citrobacter freundii</i> ATCC 8090 | 3.9 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Enterobacter aerogenes</i> ATCC 13048 | 2.35 X 10 ⁷ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Enterobacter agglomerans</i> ATCC 27155 | 3.9 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Enterobacter cloacae</i> ATCC 13047 | 3.3 X 10 ⁷ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Enterococcus faecalis</i> ATCC 19433 | 6.2 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Enterococcus faecalis</i> Vancomycin Resistant (VRE) ATCC 51299 | 1.3 X 10 ⁷ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |

AIRX 78PLUS[®]

Disinfectant Cleaner
& Odor Counteractant

Efficacy
Data

EPA REG. # 10324-155-44089

| ORGANISM | CARRIER POPULATION | SAMPLE | # CARRIERS | # POSITIVE |
|---|------------------------------------|--------|------------|------------|
| <i>Enterococcus hirae</i> ATCC 10541 | 1.19 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Escherichia coli</i> ATCC 11229 | 1.3 X 10 ⁷ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Escherichia coli</i> (carbapenem resistant) CDC 87371 | 1.34 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | 7.2 X 10 ⁵ CFU/Carrier | B | 10 | 0/10 |
| <i>Escherichia coli</i> O111 :H8 ATCC BAA-184 | 4.3 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Escherichia coli</i> Tetracycline Resistant ATCC 47041 | 3.1 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Escherichia coli</i> Spectrum B-Lactamase (ESBL) ATCC BAA-196 | 4.6 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Fusobacterium necrophorum</i> ATCC 27852 | 5.8 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Klebsiella oxytoca</i> ATCC 13182 | 1.07 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Klebsiella pneumoniae</i> ATCC 13883 | 1.2 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Klebsiella pneumoniae</i> (New Delhi metallo-beta-lactamase) CDC 1000527 | 9.5 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Listeria monocytogenes</i> ATCC 19117 | 7.7 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Micrococcus luteus</i> ATCC 14452 | 1.1 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Pasturella multocida</i> ATCC 12947 | 1.32 X 10 ⁷ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Proteus vulgaris</i> ATCC 13315 | 1.9 X 10 ⁴ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Pseudomonas aeruginosa</i> Tetracycline Resistant ATCC 27853 | 3.5 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Pseudomonas cepacia</i> ATCC 25416 | 1.63 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Salmonella enterica</i> ATCC 23564 | 9.2 X 10 ⁴ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Salmonella enteritidis</i> ATCC 4931 | 1.52 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Salmonella enterica</i> serotype pullorum ATCC 19945 | 7.1 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Salmonella typhi</i> ATCC 6539 | 8.3 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Salmonella typhimurium</i> ATCC 23564 | 1.5 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | 5.6 X 10 ⁶ CFU/Carrier | B | 10 | 0/10 |

AIRX 78PLUS[®]

Disinfectant Cleaner
& Odor Counteractant

Efficacy
Data

EPA REG. # 10324-155-44089

| ORGANISM | CARRIER POPULATION | SAMPLE | # CARRIERS | # POSITIVE |
|---|------------------------------------|--------|------------|------------|
| <i>Serratia marcescens</i> ATCC 14756 | 6.2 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Shigella flexneri</i> ATCC 9380 | 1.99 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Shigella sonnei</i> ATCC 25931 | 1.04 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus aureus</i> ATCC 25923 | 6.6 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus aureus sub species aureus</i> ATCC 33586 | 7.2 X 10 ⁴ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus aureus Methicillin Resistant (MRSA)</i> ATCC 33592 | 5.4 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus aureus Community Associated Methicillin Resistant (CA-MRSA)</i> | 6.3 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus aureus Vancomycin Intermediate Resistant (VISA)</i> ATCC 5836 | 3.2 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus epidermidis</i> ATCC 14990 | 1.56 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus epidermidis Antibiotic resistant</i> ATCC 51625 | 8.6 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Staphylococcus haemolyticus</i> ATCC 29970 | 9.5 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Streptococcus agalactiae</i> ATCC 13813 | 5.6 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Streptococcus mutans</i> ATCC 25175 | 1.02 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | 1.3 X 10 ⁴ CFU/Carrier | B | 10 | 0/10 |
| <i>Streptococcus pneumoniae Penicillin Resistant</i> ATCC 51915 | 9.6 X 10 ⁴ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Streptococcus pyogenes</i> ATCC 19615 | 4.7 X 10 ⁴ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Vibrio cholera</i> ATCC 11623 | 1.0 X 10 ⁶ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| <i>Yersinia enterocolitica</i> ATCC 23715 | 1.2 X 10 ⁷ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |

(Testing is performed per EPA Guidance (DIS/TSS-7). Two separate lots are tested. Inactivation of virus must be demonstrated at all dilutions when no cytotoxicity is observed or at all dilutions above the cytotoxic level when it is observed. The data must demonstrate a 3-log reduction in viral titer for both lots.)

| ORGANISM | DRIED VIRUS CONTROL | SAMPLE | RESULT | LOG REDUCTION |
|--|------------------------|--------|------------------------|-------------------------|
| <i>Chlamydia psittaci</i> ATCC VR-125 | 7.25 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥6.75 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥6.75 Log ₁₀ |
| | 4.75 Log ₁₀ | C | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |

AIRX[®] 78 PLUS

Disinfectant Cleaner
& Odor Counteractant

Efficacy
Data

EPA REG. # 10324-155-44089

Virucidal against (at 1 ounce per gallon)

AIRX78 PLUS was evaluated in the presence of 5% serum and 400 ppm hard water with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces.

(Testing is performed per EPA Guidance (DIS/TSS-7). Two separate lots are tested. Inactivation of virus must be demonstrated at all dilutions when no cytotoxicity is observed or at all dilutions above the cytotoxic level when it is observed. The data must demonstrate a 3-log reduction in viral titer for both lots.) (3 lots and 4-Log reduction for Canada).

| ORGANISM | DRIED VIRUS CONTROL | SAMPLE | RESULT | LOG REDUCTION |
|---|------------------------|----------------|------------------------|-------------------------|
| Avian Infectious Bronchitis virus Beaudette IB42 | 6.42 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.92 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.92 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥6.0 Log ₁₀ |
| Avian Influenza A (H3N2) virus (Avian Reassortant) (ATCC VR-2072) | 4.75 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| Avian Influenza A (H5N1) virus | 6.75 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥6.25 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥6.25 Log ₁₀ |
| Canine Coronavirus ATCC VR-809 | 4.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| Canine Distemper virus ATCC VR-128 | 6.25 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.75 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.75 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥6.25 Log ₁₀ |
| SARS Associated Coronavirus | 6.5 Log ₁₀ | A | ≤3.5 Log ₁₀ | ≥3.0 Log ₁₀ |
| | | B | ≤3.5 Log ₁₀ | ≥3.0 Log ₁₀ |
| Cytomegalovirus ATCC VR-538 | 4.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| Feline Picornavirus ATCC VR-649 | 4.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥5.25 Log ₁₀ |
| Hepatitis B Virus | 5.06 Log ₁₀ | A | 0.27 Log ₁₀ | 4.79 Log ₁₀ |
| | | B | 0.41 Log ₁₀ | 4.79 Log ₁₀ |
| | | Confirmatory B | 0.27 Log ₁₀ | 4.79 Log ₁₀ |
| Hepatitis C Virus | 6.21 Log ₁₀ | A | 0.24 Log ₁₀ | 5.97 Log ₁₀ |
| | | B | 0.42 Log ₁₀ | 5.79 Log ₁₀ |
| | | Confirmatory B | 0.13 Log ₁₀ | 5.93 Log ₁₀ |

AIRX 78PLUS[®]

Disinfectant Cleaner
& Odor Counteractant

Efficacy
Data

EPA REG. # 10324-155-44089

| | DRIED VIRUS CONTROL | SAMPLE | RESULT | LOG REDUCTION |
|---|------------------------|--------|------------------------|-------------------------|
| Herpes Simplex Virus Type 1 ATCC VR-773 | 5.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥5.5 Log ₁₀ |
| Herpes Simplex Virus Type 2 ATCC VR-734 | 6.0 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.5 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.5 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.5 Log ₁₀ |
| Human Coronavirus ATCC VR-740 | 4.5 Log ₁₀ | A | ≥0.5 Log ₁₀ | ≤4.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| Human Immunodeficiency Virus type 1 (HIV 1) HTLV-III _B | 5.75 Log ₁₀ | A | ≤1.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| | | B | ≤1.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| | | C | ≤1.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| Infectious Bovine Rhinotracheitis virus ATCC VR-188 | 4.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| Influenza A virus (H3N2) (Hong King Strain) ATCC VR-544 | 6.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥6.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥6.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥5.5 Log ₁₀ |
| Influenza A (H1N1) virus ATCC VR-1469 | 5.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |
| Pseudorabies virus ATCC VR-135 | 6.25 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.75 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.75 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |
| Respiratory syncytial virus ATCC VR-26 | 4.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥4.5 Log ₁₀ |
| Swine Influenza A (H1N1) Virus ATCC VR-333 | 5.5 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥5.0 Log ₁₀ |

AIRX[®] 78 PLUS

Disinfectant Cleaner
& Odor Counteractant

**Efficacy
Data**

EPA REG. # 10324-155-44089

| | | | | |
|-------------------------------------|------------------------|---|------------------------|-------------------------|
| Transmissible Gastroenteritis virus | 4.75 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥4.25 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥5.75 Log ₁₀ |
| Vaccinia virus ATCC VR-119 | 6.75 Log ₁₀ | A | ≤0.5 Log ₁₀ | ≥6.25 Log ₁₀ |
| | | B | ≤0.5 Log ₁₀ | ≥6.25 Log ₁₀ |
| | | C | ≤0.5 Log ₁₀ | ≥6.0 Log ₁₀ |

Virucidal against (at 4 ounces per gallon)

AIRX78 PLUS was evaluated in the presence of 5% serum and 400 ppm hard water with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces.

(Testing is performed per EPA Guidance (DIS/TSS-7). Two separate lots are tested. Inactivation of virus must be demonstrated at all dilutions when no cytotoxicity is observed or at all dilutions above the cytotoxic level when it is observed. The data must demonstrate a 3-log reduction in viral titer for both lots.) (3 lots and 4-Log reduction for Canada).

| ORGANISM | DRIED VIRUS CONTROL | SAMPLE | RESULT | LOG REDUCTION |
|--|------------------------|--------|------------------------|-------------------------|
| Canine Parvovirus Type 2b, Nike Strain | 7.5 Log ₁₀ | A | ≤3.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| | | B | ≤3.5 Log ₁₀ | ≥4.0 Log ₁₀ |
| Rabies Virus | 5.75 Log ₁₀ | A | ≤2.5 Log ₁₀ | ≥3.25 Log ₁₀ |
| | | B | ≤2.5 Log ₁₀ | ≥3.25 Log ₁₀ |
| Porcine Parvovirus (VR-742) | 5.5 Log ₁₀ | A | ≤2.5 Log ₁₀ | ≥3.0 Log ₁₀ |

Fungicidal against (at 1 ounce per gallon)

AIRX78 PLUS was evaluated in the presence of 5% serum and 400 ppm hard water with a 10 minute contact time and found to be effective against the following fungi on hard nonporous environmental surfaces.

(Testing is performed per the AOAC fungicidal method (DIS/TSS-6). Two separate lots are tested against Trichophyton mentagrophytes in a suspension test. Killing of all fungal spores in 10 minutes is required.)

| ORGANISM | CARRIER POPULATION | SAMPLE | # CARRIERS | # POSITIVE |
|--|------------------------------------|--------|------------|------------|
| Candida albicans ATCC #10231 | 1.57 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |
| Trichophyton mentagrophytes ATCC #9533 | 1.10 X 10 ⁵ CFU/Carrier | A | 10 | 0/10 |
| | | B | 10 | 0/10 |

Mold and Mildew Control (at 1 ounce per gallon)

Use **AIRX78 PLUS** to control the growth of mold and mildew and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

| ORGANISM | TILE NUMBER | UNTREATED AFTER 7 DAYS | SAMPLE A AFTER 7 DAYS | SAMPLE B AFTER 7 DAYS |
|-------------------------------|-------------|------------------------|-----------------------|-----------------------|
| Aspergillus niger ATCC #16404 | 1 | Growth 90% | No Growth 0% | No Growth 0% |
| | 2 | Growth 70% | No Growth 0% | No Growth 0% |
| | 3 | Growth 90% | No Growth 0% | No Growth 0% |